

Valence-bond crystalline order in the $s = 1/2$ J_1 - J_2 model on the honeycomb lattice

R. F. Bishop,¹ P. H. Y. Li,¹ and C. E. Campbell²

¹*School of Physics and Astronomy,*

The University of Manchester, Manchester, M13 9PL, UK

²*School of Physics and Astronomy, University of Minnesota,*

Minneapolis, Minnesota 55455, USA

Abstract

Using the coupled cluster method we study the phase diagram of the spin-1/2 Heisenberg antiferromagnet on a honeycomb lattice with nearest-neighbor exchange coupling $J_1 > 0$ and frustrating next-nearest-neighbor coupling $J_2 \equiv xJ_1 > 0$. In the range $0 < x < 1$ we find four phases exhibiting respectively Néel, plaquette, staggered dimer, and Néel-II orderings, with quantum critical points at $x_{c1} \approx 0.207(3)$, $x_{c2} \approx 0.385(10)$, and $x_{c3} \approx 0.65(5)$. The transitions at x_{c1} and x_{c3} appear to be continuous (and hence deconfined) ones, while that at x_{c2} appears to be a direct first-order one.

PACS numbers: 75.10.Jm, 75.10.Kt, 75.30.Kz, 75.50.Ee

Frustrated quantum spin models on the two-dimensional (2D) honeycomb lattice have become the objects of intense study. Impetus came from the reported presence of a quantum spin-liquid (QSL) phase in both the exactly soluble (albeit somewhat artificial) Kitaev model of spin-1/2 particles on a honeycomb lattice [1], and the half-filled Fermi-Hubbard (FH) model on a honeycomb lattice [2]. Thus, Meng *et al.* reported in a quantum Monte Carlo (QMC) calculation, free of the usual fermion sign problems, the presence in the honeycomb FH model of a QSL phase, at moderate values of the on-site Coulomb repulsion strength (U), situated between the nonmagnetic metallic insulator (or semi-metal) phase at low U and the antiferromagnetic (AFM) Mott insulator phase for large U . Since the $U \rightarrow \infty$ limit corresponds to the pure Heisenberg antiferromagnet (HAFM), i.e., with nearest-neighbor (NN) interactions (of strength $J_1 > 0$) only, the Mott insulator phase of the Hubbard model corresponds to the Néel-ordered phase of the HAFM spin-lattice model. Higher-order terms in the t/U expansion of the FH model (where t is the strength parameter of the NN hopping term) lead to frustrating exchange couplings in the corresponding spin-lattice model in which the HAFM with NN exchange couplings is the leading term in the large- U expansion. The simplest such frustrated model is the J_1 - J_2 model studied here, where the next-nearest-neighbor (NNN) spin pairs have an additional exchange coupling of strength $J_2 > 0$.

A later study of the FH model, using a Schwinger boson mean field theory (SB-MFT) approach [3], provided some corroborating evidence for a $\mathbb{Z}_2$ QSL state; and a Schwinger fermion representation of the same model [4] gave some evidence for both a $\mathbb{Z}_2$ QSL phase and a chiral antiferromagnetic phase. However, later numerically exact QMC calculations by Sorella *et al.* [5], with much larger clusters than those used by Meng *et al.* [2], have cast considerable doubt on their original finding of an intermediate QSL phase. Nevertheless, the study of analogous frustrated spin models on the honeycomb lattice remains of great interest in its own right. Thus, at intermediate values of U , around those for which Meng *et al.* [2] initially reported the QSL, the properties of the FH model are mirrored, at least loosely, by a spin-1/2 J_1 - J_2 model on the honeycomb lattice. For this and other reasons, this model and its generalizations [specifically to include also next-next-nearest-neighbor (NNNN) bonds with strength J_3], have been much studied [6–16] recently. Its Hamiltonian




FIG. 1: (Color online) The J_1 - J_2 model on the honeycomb lattice (with $J_1 = 1$), showing (a) the Néel and (b) Néel-II states. The arrows represent spins located on lattice sites $\bullet$.

is given by

$$H = J_1 \sum_{\langle i,j \rangle} \mathbf{s}_i \cdot \mathbf{s}_j + J_2 \sum_{\langle\langle i,k \rangle\rangle} \mathbf{s}_i \cdot \mathbf{s}_k, \quad (1)$$

where index i runs over all honeycomb lattice sites, and indices j and k run over all NN and NNN sites to i , respectively, counting each bond once only. Each lattice site i carries a particle with spin $s = \frac{1}{2}$ and a spin operator $\mathbf{s}_i = (s_i^x, s_i^y, s_i^z)$. The lattice and exchange bonds are illustrated in Fig. 1. We are interested in the case where both NN and NNN bonds are AFM in nature, and henceforth, we put $J_1 = 1$ to set the energy scale and define the frustration parameter $x \equiv J_2/J_1$.

The classical ($s \rightarrow \infty$) ground state of the model is Néel-ordered for $0 \leq x < \frac{1}{6}$, whereas for all values $x > \frac{1}{6}$ the spins are spirally ordered. In this latter regime, the classical model has a one-parameter family of degenerate incommensurate ground states where the spiral wave vector can orient in any direction. At leading order, i.e., $O(1/s)$, spin-wave fluctuations lift this accidental degeneracy in favor of particular wave vectors [6]. For the extreme quantum case, $s = 1/2$, considered here, we expect quantum fluctuations to be strong enough to destroy the spiral order over a wide range of values of x . In a recent paper [12] that used the coupled cluster method (CCM), we have verified that expectation for all values in the range $0 \leq x \leq 1$ considered here.

We showed too [12] that quantum fluctuations preserve the Néel order to higher values of x than in the classical model. Thus, we found that the ground-state (GS) phase of the $s = 1/2$ model is Néel-ordered for $x < x_{c1} \approx 0.207(3)$. At $x = x_{c1}$ there appears to be a continuous deconfined phase transition to a GS paramagnetic phase exhibiting plaquette valence-bond crystalline (PVBC) order. Furthermore, we found the PVBC state to be the

stable GS phase in the regime $x_{c_1} < x < x_{c_2}$, where $x_{c_2} \approx 0.385(10)$. Our aim now is to investigate further the transition at $x = x_{c_2}$ and the nature of the GS phase(s) for $x > x_{c_2}$.

The CCM, that we will employ here, has been very successfully applied to many models in quantum magnetism, including models on the honeycomb lattice [7, 12, 13] of interest here. It provides a well-structured means of studying various candidate GS phases and their regimes of stability, for each of which the description is systematically improvable in terms of well-defined truncation hierarchies for the quantum multi-spin correlations. We now briefly describe the method and refer the reader to Refs. [17–23] for further details.

The starting point for any CCM calculation is the selection of a suitable normalized model (or reference) state $|\Phi\rangle$. For spin systems it is often convenient to take a classical (uncorrelated) GS wave function for $|\Phi\rangle$. For the present case we choose the Néel state shown in Fig. 1(a) for small values of the frustration parameter x . For larger values of x we could choose one of the classical spiral GS phases to provide a CCM model state, but as we have argued above these are likely to be very fragile against quantum fluctuations. Instead, for larger values of x , we choose here the so-called Néel-II phase shown in Fig. 1(b) (which has also been denoted as the anti-Néel phase earlier [12]), that occurs in the classical ($s \rightarrow \infty$) model only at the isolated and highly degenerate critical point $x = \frac{1}{2}$. Whereas the Néel state has all 3 NN spins to a given spin antiparallel to it, the Néel-II state also comprises AFM sawtooth chains along one of the three equivalent honeycomb directions, but with NN spins on adjacent chains now parallel to one another. The Néel-II state thus breaks the lattice rotational symmetry.

It is convenient to perform a mathematical rotation of the local axes of the spins such that all spins in the reference state align along the negative z -axis. The Schrödinger ground-state ket and bra CCM equations are $H|\Psi\rangle = E|\Psi\rangle$ and $\langle\tilde{\Psi}|H = E\langle\tilde{\Psi}|$ respectively. The CCM employs the exponential parametrizations, $|\Psi\rangle = e^S|\Phi\rangle$ and $\langle\tilde{\Psi}| = \langle\Phi|\tilde{S}e^{-S}$. The correlation operator S is expressed as $S = \sum_{I \neq 0} \mathcal{S}_I C_I^+$ and its counterpart is $\tilde{S} = 1 + \sum_{I \neq 0} \tilde{\mathcal{S}}_I C_I^-$ where, by definition, $C_I^-|\Phi\rangle = 0 = \langle\Phi|C_I^+, \forall I \neq 0$. Thus we have the normalization condition $\langle\tilde{\Psi}|\Psi\rangle = \langle\Phi|\Phi\rangle \equiv 1$. The multispin creation operators $C_I^+ \equiv (C_I^-)^\dagger$, with $C_0^+ \equiv 1$, are written as $C_I^+ \equiv s_{j_1}^+ s_{j_2}^+ \cdots s_{j_n}^+$, in terms of the single-site spin-raising operators $s_k^+ \equiv s_k^x + i s_k^y$. The GS energy is $E = \langle\Phi|e^{-S} H e^S|\Phi\rangle$; and the local average onsite magnetization M in the rotated spin coordinates is $M \equiv -\frac{1}{N} \langle\tilde{\Psi}|\sum_{j=1}^N s_j^z|\Psi\rangle$. The ket- and bra-state correlation coefficients $(\mathcal{S}_I, \tilde{\mathcal{S}}_I)$ are calculated by requiring the expectation value $\bar{H} = \langle\tilde{\Psi}|H|\Psi\rangle$ to be

a minimum with respect to all parameters $(\mathcal{S}_I, \tilde{\mathcal{S}}_I)$, and hence $\langle \Phi | C_I^- e^{-S} H e^S | \Phi \rangle = 0$ and $\langle \Phi | \tilde{S}(e^{-S} H e^S - E_0) C_I^+ | \Phi \rangle = 0 ; \forall I \neq 0$.

The CCM formalism is exact if all spin configurations are included in the S and $\tilde{S}$ operators. In practice, however, truncations are needed. We employ here the well-studied localized (lattice-animal-based subsystem) LSUB m scheme [18–23], in which all possible multi-spin-flip correlations over different locales on the lattice defined by m or fewer contiguous lattice sites are retained. Such clusters are defined to be contiguous in this sense if every site in the cluster is adjacent (as a nearest neighbor) to at least one other site in the cluster. The numbers N_f of such fundamental configurations that are distinct under the (space and point-group) symmetries of the lattice and the model state increase rapidly with the LSUB m truncation index m . Thus the highest LSUB m level that we can reach here, even with massive parallelization and the use of supercomputing resources [24], is LSUB12, for which $N_f = 293309$ for the Néel-II state.

Since, in any truncation, CCM parametrizations automatically satisfy the Goldstone linked cluster theorem, we may work from the outset in the thermodynamic limit, $N \rightarrow \infty$. Nevertheless, the raw LSUB m data still need to be extrapolated to the exact $m \rightarrow \infty$ limit. Although there are no known exact extrapolation rules we have much experience in doing so. Thus, for the GS energy per spin, E/N , we use (see, e.g., Refs. [20–23])

$$E(m)/N = a_0 + a_1 m^{-2} + a_2 m^{-4}; \quad (2)$$

while for the magnetic order parameter, M , defined above, we use either the scheme

$$M(m) = b_0 + b_1 m^{-1} + b_2 m^{-2}, \quad (3)$$

for systems showing no or only slight frustration (see, e.g., Refs. [19, 21]), or the scheme

$$M(m) = c_0 + c_1 m^{-1/2} + b_2 m^{-3/2}, \quad (4)$$

for more strongly frustrated systems or ones showing a GS order-disorder transition (see, e.g., Refs. [22, 23]). Since the hexagon is an important structural element of the honeycomb lattice we perform the extrapolations here only for LSUB m data with $m \geq 6$.

In Fig. 2 we show our results for the magnetic order parameter, M , using both the Néel and Néel-II states as CCM model states. On the Néel side we show extrapolations using both Eqs. (3) and (4). Equation (3), which is appropriate when $J_2 = 0$, yields




FIG. 2: (Color online) CCM LSUB m results for the GS order parameter, M , of the spin-1/2 J_1 - J_2 honeycomb model ($J_1 = 1$), using the Néel (left curves) and Néel-II (right curves) states as model states, with $m = \{6, 8, 10, 12\}$. The extrapolated curves LSUB ∞ (1) and LSUB ∞ (3) use this data set with Eqs. (4) and (3) respectively, while the LSUB ∞ (2) curve uses Eq. (4) with the restricted set $m = \{8, 10, 12\}$.

the value $M \approx 0.271(2)$ for the unfrustrated HAFM on the hexagonal lattice (i.e., with NN interactions only), in excellent agreement with the best available QMC estimate [25], $M = 0.2677(6)$. Our own error estimates are based on sensitivity checks using different LSUB m data sets. Similarly we see from Fig. 2 that all extrapolations give essentially the same estimate $x_{c_1} \approx 0.207(3)$ for the point where Néel order vanishes ($M \rightarrow 0$). We showed previously [12] that the phase transition at $x = x_{c_1}$ is a continuous deconfined one between states with Néel and PVBC order.

Figure 2 also shows corresponding results for M for a possible phase with Néel-II ordering. Inspection shows clearly that inclusion of the LSUB m result with $m = 6$ for this state is not compatible with the expected extrapolation scheme of Eq. (4), whereas those with $m > 6$ fit very well. Why the $m = 6$ result should be anomalous in this case is unclear. For these reasons we show in Fig. 2 only extrapolated results using the Néel-II model state based on $m = \{8, 10, 12\}$. The results clearly show that Néel-II ordering is present, albeit with a rather small value of the order parameter, $M \lesssim 0.1$, for $x > x_{c_3}$ where $x_{c_3} \approx 0.65(5)$, but where the error estimate is now very rough and rather uncertain.

In our previous work [12] we showed that the Néel-II state becomes susceptible to PVBC ordering for $x < x_{c_2} \approx 0.385(10)$, but we now observe that the Néel-II state is itself only




FIG. 3: (Color online) Left: CCM LSUB m results for the inverse staggered dimer susceptibility, $1/\chi_d$, of the spin-1/2 J_1 - J_2 honeycomb model ($J_1 = 1$), using the Néel-II state as model state, with $m = \{6, 8, 10, 12\}$. The extrapolated curves LSUB ∞ (1) and LSUB ∞ (2) are derived from fitting the perturbed energies (see text) as $e(\delta) = e_0(\delta) + e_1(\delta)m^{-\nu}$, and use the data sets $m = \{6, 8, 10, 12\}$ and $m = \{8, 10, 12\}$ respectively. Right: The field $F \rightarrow \delta \hat{O}_d$ for the staggered dimer susceptibility, χ_d . Thick (red) and thin (black) lines correspond respectively to strengthened and unaltered NN exchange couplings, where $\hat{O}_d = \sum_{\langle i,j \rangle} a_{ij} \mathbf{s}_i \cdot \mathbf{s}_j$, and the sum runs over all NN bonds, with $a_{ij} = +1$ and 0 for thick (red) lines and thin (black) lines respectively.

stable as a magnetically ordered state for $x > x_{c3}$. We are thus led to enquire about the possible GS phase(s) of the system in the range $x_{c2} < x < x_{c3}$. In view of the persistence of our CCM LSUB m solutions based on the Néel-II model state, with finite values of m , well into the region $x < x_{c3}$ before they terminate (as is clearly seen from Fig. 2), we expect that the actual GS phase in this intermediate regime might share similarities with the Néel-II state. For example, just as the Néel-II state breaks the lattice rotational symmetry, so does another valence-bond solid state, namely the staggered-dimer valence-bond crystalline (SDVBC) (or lattice nematic) state. This is formed from the Néel-II state by replacing all of the parallel NN spin pairs by spin-zero dimers (and see Fig. 3).

In order to investigate the possibility of an SDVBC phase we first consider the response of the system to a field operator F (and see, e.g., Ref. [23]). Thus, a field term $F = \delta \hat{O}_d$ is added to the Hamiltonian of Eq. (1), where $\hat{O}_d$ is an operator corresponding to the possible SDVBC order, illustrated in Fig. 3 and defined in its caption. The energy per site, $E(\delta)/N \equiv$

$e(\delta)$, is then calculated in the CCM for the perturbed Hamiltonian $H + F$, using the Néel-II model state. We define the corresponding susceptibility as $\chi_d \equiv -(\partial^2 e(\delta))/(\partial \delta^2)|_{\delta=0}$. Clearly the GS phase becomes unstable against SDVBC order when χ_d^{-1} becomes zero. We now use the LSUB m extrapolation scheme $e(\delta) = e_0(\delta) + e_1(\delta)m^{-\nu}$, with the exponent ν also a fitting parameter, rather than our standard energy extrapolation scheme of Eq. (2), in order to calculate the extrapolated values of χ_d^{-1} shown in Fig. 3. For the same data set $m = \{8, 10, 12\}$ used to calculate M for the Néel-II state above, the fitted value of ν is close to 2 over most of the range of the J_2 values shown, except near the termination point of this phase, where it falls sharply. By contrast, for the set $m = \{6, 8, 10, 12\}$ also shown in Fig. 3, ν is closer to 1 over most of the range. This again reinforces the anomalous nature of the LSUB6 results.

What we see from Fig. 3 is that the extrapolated value of χ_d^{-1} is close to zero over a range of values of x that extends from x_{c_2} below to an upper value of about 0.6, which is completely compatible with the value x_{c_3} obtained from the order parameter M of the Néel-II state. Thus, by combining our results, we conclude that in the region $x_{c_2} < x < x_{c_3}$ the GS phase has SDVBC order, while for $x > x_{c_3}$ the GS phase has Néel-II order, although this latter ordering is weak and quite fragile against the still strongly competing SDVBC order. The shape of the CCM curves for χ_d^{-1} in Fig. 3 are indicative of a continuous (and hence deconfined) quantum critical point at x_{c_3} , whereas the corresponding curves for χ_p^{-1} , the inverse plaquette susceptibility, found in our earlier work [12] were much more indicative of a direct first-order transition at x_{c_2} . We see no signals at all of the spiral ordering that is present classically for $x > \frac{1}{6}$ for any value of x in the range $0 < x < 1$ examined.

In conclusion, over the range $0 < x < 1$ we find that the spin-1/2 J_1 - J_2 HAFM on the honeycomb lattice has four phases with, respectively, Néel, PVBC, SDVBC, and Néel-II ordering. Our first calculated critical point, $x_{c_1} \approx 0.207(3)$, at which Néel order melts, agrees well with other recent results, including $x_{c_1} \approx 0.195(25)$ from a large-scale exact diagonalization (ED) study [8], $x_{c_1} \approx 0.26$ [14] and $x_{c_1} \approx 0.22$ [15] from two separate density-matrix renormalization group (DMRG) studies, and $x_{c_1} \approx 0.2075$ from a SB-MFT approach [16]. Both DMRG studies [14, 15] and the ED study [8] concur with us that the transition at x_{c_1} is probably a continuous deconfined one to a PVBC state, whereas the SB-MFT study [16] indicates a transition to a QSL state. Our second calculated critical point, $x_{c_2} \approx 0.385(10)$, at which the PVBC order melts, is similarly in good agreement with the

result $x_{c_2} \approx 0.375(25)$ from the ED study [8], and the results $x_{c_2} \approx 0.36$ [14] and $x_{c_2} \approx 0.35$ [15] from the two DMRG studies. We find that the transition at x_{c_2} is probably a direct first-order one to a state with SDVBC order. Both DMRG studies [14, 15] concur that the transition at x_{c_2} is to a state with SDVBC order, although Ganesh *et al.* [15] find evidence for the surprising scenario that the transition at x_{c_2} is also of the continuous deconfined type, as at x_{c_1} . The ED study [8], by contrast, finds a first-order transition at x_{c_2} to a state that cannot be distinguished between having either SDVBC or Néel-II order. Finally, we find evidence for a third critical point at $x_{c_3} \approx 0.65(5)$ at which a continuous (and hence again deconfined) transition occurs to a state with weak Néel-II magnetic order. It is interesting to note that such a transition is also compatible with the DMRG result of Ganesh *et al.* [15], which could not rule out a melting of the SDVBC order for values $x \gtrsim 0.7$.

We thank the University of Minnesota Supercomputing Institute for the grant of supercomputing facilities for this research.

-
- [1] A. Kitaev, Ann. Phys. (NY) **321**, 2 (2006); G. Baskaran, S. Mandal, and R. Shankar, Phys. Rev. Lett. **98**, 247201 (2007); J. Chaloupka, G. Jackeli, and G. Khaliullin, *ibid.* **105**, 027204 (2010).
 - [2] Z. Y. Meng, T. C. Lang, S. Wessel, F. F. Assaad, and A. Muramatsu, Nature (London) **464**, 847 (2010).
 - [3] F. Wang, Phys. Rev. B **82**, 024419 (2010).
 - [4] Y.-M. Lu, and Y. Ran, Phys. Rev. B **84**, 024420 (2011).
 - [5] S. Sorella, Y. Otsuka, and S. Yunoki, Scientific Reports **2**, 992 (2012).
 - [6] A. Mulder, R. Ganesh, L. Capriotti, and A. Paramekanti, Phys. Rev. B **81**, 214419 (2010).
 - [7] D. J. J. Farnell, R. F. Bishop, P. H. Y. Li, J. Richter, and C. E. Campbell, Phys. Rev. B **84**, 012403 (2011).
 - [8] A. F. Albuquerque, D. Schwandt, B. Hetényi, S. Capponi, M. Mambrini, and A. M. Läuchli, Phys. Rev. B **84**, 024406 (2011).
 - [9] J. Oitmaa and R. R. P. Singh, Phys. Rev. B **84**, 094424 (2011).
 - [10] J. Reuther, D. A. Abanin, and R. Thomale, Phys. Rev. B **84**, 014417 (2011).
 - [11] F. Mezzacapo and M. Boninsegni, Phys. Rev. B **85**, 060402(R) (2012).

- [12] R. F. Bishop, P. H. Y. Li, D. J. J. Farnell, and C. E. Campbell, *J. Phys.: Condens. Matter* **24**, 236002 (2012).
- [13] P. H. Y. Li, R. F. Bishop, D. J. J. Farnell, and C. E. Campbell, *Phys. Rev. B* **86**, 144404 (2012).
- [14] Z. Zhu, D. A. Huse, and S. R. White, e-print arXiv:1212.6322v1.
- [15] R. Ganesh, J. van den Brink, and S. Nishimoto, e-print arXiv:1301.0853v1.
- [16] H. Zhang and C. A. Lamas, *Phys. Rev. B* **87**, 024415 (2013).
- [17] R. F. Bishop, *Theor. Chim. Acta* **80**, 95 (1991).
- [18] C. Zeng, D. J. J. Farnell, and R. F. Bishop, *J. Stat. Phys.* **90**, 327 (1998).
- [19] S. E. Krüger, J. Richter, J. Schulenburg, D. J. J. Farnell, and R. F. Bishop, *Phys. Rev. B* **61**, 14607 (2000).
- [20] R. F. Bishop, D. J. J. Farnell, S. E. Krüger, J. B. Parkinson, J. Richter, and C. Zeng, *J. Phys. Condens. Matter* **12**, 6887 (2000).
- [21] R. Darradi, J. Richter, and D. J. J. Farnell, *Phys. Rev. B* **72**, 104425 (2005).
- [22] R. F. Bishop, P. H. Y. Li, R. Darradi, J. Schulenburg, and J. Richter, *Phys. Rev. B* **78**, 054412 (2008).
- [23] R. Darradi, O. Derzhko, R. Zinke, J. Schulenburg, S. E. Krüger, and J. Richter, *Phys. Rev. B* **78**, 214415 (2008).
- [24] We use the program package CCCM of D.J.J. Farnell and J. Schulenburg, see <http://www-e.uni-magdeburg.de/jschulen/ccm/index.html>.
- [25] E. V. Castro, N. M. R. Peres, K. S. D. Beach, and A. W. Sandvik, *Phys. Rev. B* **73**, 054422 (2006).